Wave 2025

**JACK PHU**

DAY 1 – Session 3 – Interactive Polls using Slido with Jack Phu

Advance preview for your reference only – You will need to answer each when Slido is live during Jack's presentation. Questions only appear when Jack is on that slide.

WAVE 2025 Intera...

Poll 1. A history of trauma to the side of the head is most likely to lead to which type of optic neuropathy? 0

WAVE 2025 Polls Only Active During Interactive Session.

- A. Primary optic atrophy
- B. Secondary optic atrophy
- C. Consecutive optic atrophy
- D. Trans-synaptic retrograde degeneration
- E. Glaucoma

Send

WAVE 2025 Intera...

Poll 2. Which of the following represents best practice for reporting intraocular pressure for glaucoma? 0

- A. Report a range to capture variability
- B. Report a single, average value only
- C. Correct for corneal thickness
- D. Correct for corneal hysteresis
- E. Do not bother – IOP is useless for glaucoma

WAVE 2025 Intera...

Poll 3. A patient with glaucomatous neuroretinal rim loss at the 1 o'clock position (superotemporal) of left eye will most likely have which visual field defect? 0

- A. Superior arcuate
- B. Superior paracentral
- C. Enlarged blind spot
- D. Inferior paracentral
- E. Inferior arcuate

Send

Voting as Anonymous

WAVE 2025 Intera...

Poll 4. Which combination represents the most likely scenario for a poor prognosis for glaucoma? 0

- A. Superior arcuate
- B. Superior paracentral
- C. Enlarged blind spot
- D. Inferior paracentral
- E. Inferior arcuate

Send

Voting as Anonymous

WAVE 2025 Intera...

Poll 5. Next most appropriate management step for a patient with history of head trauma who has a homonymous right superior quadrantopia on examination? 0

- A. Nothing – history is clear
- B. OCT to confirm
- C. Perform fundus autofluorescence
- D. Repeat visual field testing – this is an atypical pattern
- E. Request neuroimaging

Send

Access via QR

Access via App

or www.slido.com and code 3450 959